arXiv:0809.3176v1 [hep-ph] 18 Sep 2008

Sneutrino LSPs in R-parity violating minimal supergravity models

Markus A. Bernhardt, Siba Prasad Das, Herbi K. Dreiner and Sebastian Grab

Physikalisches Institut der Universität Bonn, Nußallee 12, D-53115 Bonn, Germany

Abstract. We consider the minimal supergravity model (mSUGRA) with one additional R-parity violating operator at the GUT scale. The superparticles mass spectra at the weak scale are generally altered due to the presence of the R-parity violating coupling in the renormalization group equations. We show that a lepton number violating coupling at the GUT scale can lead to a sneutrino as the lightest supersymmetric particle (LSP) in a large region of parameter space consistent with the muon anomalous magnetic moment and other precision measurements. We also give characteristic collider signatures at the LHC.

Keywords: Supergravity, Renormalization group evolution of parameters, Supersymmetric models, Supersymmetric partners of known particles

PACS: 04.65.+e,11.10.Hi,12.60.Jv,14.80.Ly

INTRODUCTION

The most general gauge invariant and renormalizable superpotential of the supersymmetric standard model with minimal particle content contains lepton and baryon number violating operators [1],

$$W_{\not R_p} = \frac{1}{2}\lambda_{ijk}L_iL_j\bar{E}_k + \lambda'_{ijk}L_iQ_j\bar{D}_k + \frac{1}{2}\lambda''_{ijk}\bar{U}_i\bar{D}_j\bar{D}_k + \kappa_i L_i H_u, \qquad (1)$$

where $i, j, k = 1, 2, 3$ are the generation indices.

The fourth term in Eq. (1) can be rotated away at any scale by an appropriate field redefinitions [2]. Simultaneous presence of lepton (first two terms) and baryon (third term) number violating operators lead to rapid proton decay [3]. An additional discrete symmetry is therefore required to keep the proton stable [4]. R-parity, R_p, and proton-hexality, P_6, prohibits $W_{\not R_p}$. Here, we will consider a third possibility, baryon-triality, B_3, which violates R_p and P_6 by prohibiting only the $\bar{U}\bar{D}\bar{D}$ operator in Eq. (1).

The remaining two lepton number violating interactions in Eq. (1) will lead to the decay of the lightest supersymmetric particle (LSP). Since the LSP is not stable, any SUSY particle (sparticle) is allowed to be the LSP [5]. However, the nature of the LSP is crucial for supersymmetric signatures at colliders, since typically all heavier sparticle decay chains end up with the LSP.

In the B_3 minimal supergravity (mSUGRA) model [1], we assume an additional coupling Λ' at the GUT scale ($M_{\rm GUT}$). We thus have the six free parameters:

$$M_0\,, M_{1/2}\,, A_0\,, \tan\beta\,, {\rm sgn}(\mu) \text{ and } \Lambda' \quad , \qquad (2)$$

where Λ' is one coupling out of λ'_{ijk}. Due to the presence of one such coupling, the collider phenomenology could be very different compared to R_p conserving mSUGRA scenarios with a stable neutralino LSP. The possible changes are the following: different running of sparticle masses and couplings [1, 6, 7]; the LSP will decay; single production of sparticles [3, 8]; the decay patterns of the sparticles are altered [3, 6, 8].

It was shown in Ref. [6] that the model possess three different LSP candidates: the lightest neutralino, $\tilde\chi_1^0$, the lightest scalar tau (stau), $\tilde\tau_1$, and the sneutrino, $\tilde\nu_i$. The $\tilde\nu_i$ is special, because its mass needs to be reduced by additional B_3 contributions in the renormalization group equations (RGEs) in order to become the LSP. We will mainly focus on that and show that a large region in the B_3 mSUGRA parameter space exists, where a $\tilde\nu_i$ is the LSP consistent with experimental constraints, *e.g.*, the anomalous magnetic moment of the muon, a_μ [9] and the LEP bounds on lighter Higgs boson mass and sneutrino masses [10].

SNEUTRINO LSPS IN B_3 MSUGRA

The dominant contributions to the RGE of $m_{\tilde\nu_i}$ can be expressed as [1]:

$$\begin{aligned} 16\pi^2\frac{d(m^2_{\tilde\nu_i})}{dt} = & -\left(\frac{6}{5}g_1^2|M_1|^2 + 6g_2^2|M_2|^2 + \frac{3}{5}g_1^2 S\right) \\ & + 6\lambda'^2_{ijk}\left[(\mathbf{m_{\tilde L}})^2_{ii} + (\mathbf{m_{\tilde Q}})^2_{jj} + (\mathbf{m_{\tilde D}})^2_{kk}\right] \\ & + 6(\mathbf{h_{D^k}})^2_{ij} \end{aligned} \qquad (3)$$

with

$$(\mathbf{h_{D^k}})_{ij} \equiv \lambda'_{ijk} \times A_0 \quad \text{at} \quad M_{\rm GUT}. \qquad (4)$$

The running of $m_{\tilde\nu_i}$ is governed by two different terms. The first term in Eq. (3) is proportional to the gauge cou-



FIGURE 1. Masses of $\tilde{\chi}_1^0$, $\tilde{\tau}_1$, $\tilde{\nu}_\mu$ and $\tilde{\mu}_L$ at M_Z as a function of $\lambda'_{231}|_{\rm GUT}$ for the SPS1a mSUGRA parameters.



FIGURE 2. Mass difference, ΔM, between the LSP and NLSP. The LSP candidates in different regions are explicitly mentioned. The blackened out region corresponds to parameter points, which possess a tachyon or where the $\tilde{\nu}_\mu$ or the Higgs boson mass violates the LEP bounds of Ref. [10].

plings g_1, g_2 and gaugino mass parameters M_1, M_2. S is identically zero at $M_{\rm GUT}$ due to the universal nature of scalar masses in mSUGRA, see Ref. [1]. For vanishing λ'_{ijk}, $m_{\tilde{\nu}_i}$ increases while running from $M_{\rm GUT}$ to the electroweak scale (M_Z) due to the presence of the negative sign in front of the gauge terms.

The contribution from the second term is proportional to λ'_{ijk} and $(\mathbf{h}_{D^k})_{ij}$, which is also proportional to λ'_{ijk} at $M_{\rm GUT}$, *cf.* Eq. (4). This terms are, in contrast to the gauge terms, positive and will therefore reduce $m_{\tilde{\nu}_i}$ when we go from $M_{\rm GUT}$ to M_Z. The influence of these new contributions on $m_{\tilde{\nu}_i}$ depend on the magnitude of λ'_{ijk} and also on the other mSUGRA parameters, especially on A_0 [11].

We show in Fig. 1 the impact of a non-vanishing coupling $\lambda'_{ijk}|_{\rm GUT}$ on the running of $m_{\tilde{\nu}_i}$ for the SPS1a parameter set [12]. For this purpose we use an unpublished B_3 version of `SOFTSUSY` [13]. We observe, that $m_{\tilde{\nu}_\mu}$ is decreasing for increasing $\lambda'_{231}|_{\rm GUT}$ as described by Eq. (3). Furthermore, the mass of the left-handed smuon, $\tilde{\mu}_L$, also decreases because it belongs to the same SU(2) doublet like the $\tilde{\nu}_\mu$. The masses of the $\tilde{\chi}_1^0$ and $\tilde{\tau}_1$ are almost insensitive to $\lambda'_{231}|_{\rm GUT}$. We therefore obtain a $\tilde{\nu}_\mu$-LSP for $\lambda'_{231} > 0.12|_{\rm GUT}$. This large coupling is still consistent with experimental observations [3].

In general, there are large regions in the B_3 mSUGRA parameter space where the $\tilde{\nu}_i$ becomes the LSP, if $\lambda'_{ijk}|_{\rm GUT} \gtrsim \mathcal{O}(0.1)$. As an example we choose A_0 = -600 GeV, $\tan\beta$=10, sgn(μ) = +1, $\lambda'_{231} = 0.11|_{\rm GUT}$ and vary M_0 and $M_{1/2}$. We show in Fig. 2 the mass difference between the LSP and the next-to-LSP (NLSP) in the M_0–$M_{1/2}$ plane. We observe three different LSP candidates, namely the $\tilde{\nu}_\mu$, the $\tilde{\chi}_1^0$ and the $\tilde{\tau}_1$, in different regions of B_3 mSUGRA parameter space. Increasing $M_{1/2}$ increases the mass of the (left-handed) $\tilde{\nu}_\mu$ faster than the mass of the (mainly right-handed) $\tilde{\tau}_1$. We thus find a $\tilde{\tau}_1$-LSP instead of a $\tilde{\nu}_\mu$-LSP for large $M_{1/2}$. Increasing M_0, increases the masses of the sfermions but not the mass of the (bino-like) $\tilde{\chi}_1^0$ and we therefore reobtain the $\tilde{\chi}_1^0$-LSP. We also see that the LSP and NLSP are almost degenerate in mass around the thick black contours.

We conclude that a $\tilde{\nu}_\mu$-LSP exists in an extended region of B_3 mSUGRA parameter space, which is also consistent with the observed anomalous magnetic moment of the muon, a_μ. The contours in Fig. 2 correspond to the measured central value of a_μ (red), $\pm 1\sigma$ (green), $\pm 2\sigma$ (blue) and $\pm 3\sigma$ deviation (magenta) from the central value. The SUSY contributions to a_μ are calculated using `micrOMEGAs1.3.6` [14].

LHC PHENOMENOLOGY

We choose M_0=100 GeV and $M_{1/2}$=450 GeV from Fig. 2 as a representative point for investigating the $\tilde{\nu}_\mu$-LSP phenomenology at the LHC. We calculate the decay rates with `ISAWIG1.200` and `ISAJET7.64` [15]. This output is fed into `HERWIG` [16] to simulate events at the LHC. We show some masses and BRs, which are special for $\tilde{\nu}_\mu$-LSP scenarios in Table. 1.

The sparticle pair production cross-section at the LHC is 3.0 pb. Once produced, the sparticles might follow the decays in Table. 1. This will lead to very distinctive final states compared to mSUGRA with a stable $\tilde{\chi}_1^0$-LSP.

One of the most striking signatures of this scenario are high-p_T muons ($\approx$ 10% of events) from the direct decays of $\tilde{d}_R$ and $\tilde{t}_{1/2}$ via λ'_{231}, see Table. 1. We show

TABLE 1. Sparticle masses and branching ratios (BRs) in B_3 mSUGRA models. Masses which are affected due to λ'_{231} coupling and B_3 decays are shown in bold-face.

	mass	**channel**	**BR**	**channel**	**BR**
$\tilde{\nu}_\mu$	**124**	$\bar{b}d$	**100%**		
$\tilde{\mu}_L^-$	**147**	$W^-\bar{b}d$	**75.1%**	$\bar{c}d$	**24.9%**
$\tilde{\chi}_1^0$	184	$\tilde{\nu}_\mu^*\nu_\mu$	36.0%	$\tilde{\nu}_\mu\bar{\nu}_\mu$	36.0%
		$\tilde{\mu}_L^+\mu^-$	14.0%	$\tilde{\mu}_L^-\mu^+$	14.0%
$\tilde{t}_1$	**650**	$\tilde{\chi}_1^+b$	42.1%	$\tilde{\chi}_1^0t$	33.5%
		$\tilde{\chi}_2^0t$	13.8%	μ^+d	**10.6%**
$\tilde{d}_R$	**897**	$\nu_\mu b$	**45.3%**	μ^-t	**42.1%**
		$\tilde{\chi}_1^0d$	12.6%		



FIGURE 3. The normalized p_T distribution of the muon from the decays $\tilde{d}_R \to \mu t$ and $\tilde{t}_{1/2} \to \mu d$ (*cf.* Table. 1) at the LHC.

the normalized distribution in Fig. 3. The distribution peaks at 300 GeV and one can thus apply a very stringent selection cut on the muon p_T to isolate the SUSY signal from the SM background.

Additional distinctive signatures are (*cf.* Table. 1): not necessarily missing p_T ($\approx$ 25% of events), because the $\tilde{\nu}_\mu$-LSP and the $\tilde{\mu}_L$-NLSP will decay into SM particles; 2-5 non-*b* jets and 0-4 *b* jets, where most of the *b* jets stem from the LSP decay; high-p_T top quarks from $\tilde{d}_R$ decay; like-sign dimuon events mainly from pair production of right-handed squarks and subsequent decays into the (Majorana) $\tilde{\chi}_1^0$ and $\tilde{\mu}_L$. We will present the detailed analysis in [11].

CONCLUSIONS

We have shown that a coupling $\lambda'_{231}|_{\mathrm{GUT}} = \mathcal{O}(10^{-1})$ can lead to a sneutrino LSP in B_3 mSUGRA models (Fig. 1). We have found large regions of parameter space where the sneutrino is the LSP consistent with a_μ and other electroweak precision observables (Fig. 2). Distinctive LHC signatures are high-p_T muons (Fig. 3), high-p_T tops, like-sign dimuon events and *b* jets.

ACKNOWLEDGMENTS

We thank Ben Allanach for help with the yet unpublished B_3 version of `SOFTSUSY`. SPD acknowledges financial support from the Bundesministerium für Bildung und Forschung (BMBF) Projekt under Contract No. 05HT6PDA. SG thanks the "Deutsche Telekom Stiftung" and the "Bonn Cologne Graduate School" for financial support.

REFERENCES

1. B. C. Allanach, A. Dedes and H. K. Dreiner, Phys. Rev. D **69** (2004) 115002 [Erratum-ibid. D **72** (2005) 079902].
2. L. J. Hall and M. Suzuki, Nucl. Phys. B **231** (1984) 419.
3. R. Barbier *et al.*, Phys. Rept. **420** (2005) 1.
4. L. E. Ibanez and G. G. Ross, Phys. Lett. B **260** (1991) 291; L. E. Ibanez and G. G. Ross, Nucl. Phys. B **368** (1992) 3, H. K. Dreiner, C. Luhn and M. Thormeier, Phys. Rev. D **73** (2006) 075007.
5. J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos, K. A. Olive and M. Srednicki, Nucl. Phys. B **238** (1984) 453.
6. B. C. Allanach, M. A. Bernhardt, H. K. Dreiner, C. H. Kom and P. Richardson, Phys. Rev. D **75** (2007) 035002.
7. I. Jack, D. R. T. Jones and A. F. Kord, Phys. Lett. B **632** (2006) 703.
8. H. K. Dreiner, S. Grab and M. K. Trenkel, arXiv:0808.3079 [hep-ph].
9. D. Stockinger, arXiv:0710.2429 [hep-ph].
10. R. Barate *et al.*, Phys. Lett. B **565** (2003) 61; G. Abbiendi *et al.* [OPAL Collaboration], Eur. Phys. J. C **33** (2004) 149; A. Heister *et al.* [ALEPH Collaboration], Eur. Phys. J. C **31** (2003) 1.
11. M. A. Bernhardt, S. P. Das, H. K. Dreiner and S. Grab, paper in preparation.
12. B. C. Allanach *et al.*, arXiv:hep-ph/0202233.
13. B.C. Allanach and M.A. Bernhardt, unpublished.
14. G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. **149** (2002) 103.
15. F. E. Paige, S. D. Protopopescu, H. Baer and X. Tata, arXiv:hep-ph/0312045.
16. G. Corcella *et al.*, JHEP **0101** (2001) 010; G. Corcella *et al.*, S. Moretti, K. Odagiri, P. Richardson, M. H. Seymour and B. R. Webber, JHEP **0204** (2002) 028.